

# Gastropancreatic fistula in a patient with chronic pancreatitis and IPMN

Alessandro Morotti,<sup>1</sup> Marco Busso,<sup>2</sup> Paolo Rappa Verona,<sup>3</sup> Angelo Guerrasio<sup>1</sup>

<sup>1</sup>Department of Clinical and Biological Sciences, University of Turin, Orbassano, Torino, Italy

<sup>2</sup>Department of Oncology—Radiology Unit, University of Turin, Orbassano, Torino, Italy

<sup>3</sup>University of Turin, Orbassano, Italy

## Correspondence to


Dr Alessandro Morotti,  
alessandro.morotti@unito.it

AM and MB contributed equally to the manuscript.

Accepted 5 May 2016

## DESCRIPTION

A 66-year-old man was admitted to our division of internal medicine because of enteritis and sepsis. In the past, the patient had presented with chronic pancreatitis due to alcohol assumption, diabetes, cirrhosis and an intraductal papillary mucinous neoplasm (IPMN) of the pancreas (main type). At the present clinical examination, he was vomiting and had diarrhoea. Blood tests revealed leukocytosis (white cell count=19 900/ $\mu$ L), elevated PCR (27 mg/dL) and glicometabolic failure (glucose 256 mg/dL). The patient received broad-spectrum antibiotics, parenteral nutrition and rehydration, and showed sudden improvement. However, on realimentation, he developed a rapid increase of amylase (192 U/L) and lipase levels (1053 U/L), but had no abdominal pain. An abdominal sonography was performed, showing marked dilatation (17 mm) of the Wirsung duct. A CT scan of the abdomen confirmed the severe dilatation of the Wirsung duct, while the dimensions of the known IPMN were unchanged. Furthermore, a gastropancreatic fistula was observed (15 mm of diameter), connecting the pancreas body to the antral region of the stomach (figures 1 and 2). Alimentation was stopped and endoscopic examination was performed, confirming the presence of a gastric fistula. About 50% of internal and 70% of external pancreatic fistulas can be managed without the need for interventions.<sup>1</sup> However, while stenting during endoscopic retrograde cholangiopancreatography may treat fistulae associated with stenosis and gallstones, the presence of a severe (15 mm diameter) gastric communication responds to few therapeutic alternatives other than surgery. This patient underwent surgical duodenal-pancreatectomy with splenectomy and en bloc gastrectomy.



**Figure 1** CT scan showing gastropancreatic fistula.



**Figure 2** Minimum intensity projection multiplanar CT volumetric rendering with artificial green colour to highlight the gastropancreatic fistula and the Wirsung duct dilatation.

## Learning points

- ▶ Gastropancreatic fistula is a rare complication of chronic pancreatitis and can be associated with intraductal papillary mucinous neoplasm (IPMN).<sup>2,3</sup>
- ▶ Gastropancreatic fistulae can also be asymptomatic and discovered incidentally.
- ▶ Closed abdominal sonography follow-up is advisable in those patients with both chronic pancreatitis and IPMN.

**Acknowledgements** The authors would like to thank all the members of the Internal Medicine Division—San Luigi Hospital, and Professor A Veltri from the Radiology Unit of the Department of Oncology.

**Contributors** AM wrote the manuscript. AM and PRV managed the patient. MB provided CT scan images; AG reviewed clinical assessment of the patient.

**Competing interests** None declared.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 Voss M, Pappas T. Pancreatic Fistula. *Curr Treat Options Gastroenterol* 2002;5:345–53.
- 2 Honda K, Kume K, Yamasaki M, et al. Pancreatico-gastric fistulas due to intraductal papillary mucinous neoplasm (IPMN). *Intern Med* 2008;47:557–8.
- 3 Hong MY, Yu DW, Hong SG. Intraductal papillary mucinous neoplasm of the bile duct with gastric and duodenal fistulas. *World J Gastrointest Endosc* 2014;6:328–33.



CrossMark

**To cite:** Morotti A, Busso M, Rappa Verona P, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-215375

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow